Breastfeeding Associated with Lower Rates of Obesity and Lower Risk of Insulin Resistance

Breast milk feeding, compared with infant formula feeding, and breastfeeding duration are associated with lower rates of **obesity** and lower risk of **insulin resistance**.

General Information		
Broad Focus Area	Obesity and altered physical development	
Background and Justification	Although the relation of breastfeeding to subsequent overweight has been examined numerous times, only the results of two large, recent studies have provided evidence that there may be a protective effect. Data remain sparse on whether the protective association with overweight persists later in adolescence. Questions remain whether the protective effect, if any, is due to constituents of breast milk, or due to differences in feeding practices for breast and formula-fed infants. Furthermore, existing studies rely on retrospectively-reported breastfeeding histories. If breastfeeding does help prevent subsequent obesity, this would be one of the few protective preventive measures available. The relationship of large racial and ethnic differences in infant feeding practices (e.g., approximately 34% of white infants are breastfed at 6 months of age, compared to 21% of black children and 28% of Hispanic children) to subsequent differences in childhood obesity may also be explored. Identifying whether the protective effect is due to constituents of breast milk or to differences in feeding practices would help inform recommendations regarding mode of breast milk feeding in infants.	
Prevalence/ Incidence	In 2000, almost 70% of all infants were breastfed (either exclusively or in combination with formula feeding) before initial hospital discharge. However, by 6 months of age, just under one-third of infants were breastfed. ⁵ The prevalence of overweight among children is greater than 16% among children aged 6 years or more, and this prevalence has increased over the past 40 years. ^{6,7} Being overweight as a child is a risk factor for being overweight in adulthood, ⁸ and is associated with increased risk of type 2 diabetes, hypertension, and coronary artery disease. ⁹ Furthermore, being overweight as a child increases the risk of developing type 2 diabetes before the age of 21 years. ¹⁰	
Economic Impact	Because child overweight is a risk factor for adult overweight, child overweight contributes to the more than \$40 billion annual cost of obesity in the U.S. 11	

Exposure Measures		Outcome Measures	
Primary/ Maternal	Breast milk feeding and duration	Primary/ Maternal	
Methods	Interview	Methods	
Life Stage	Birth & and periodically in the first year of life	Life Stage	
Primary/Child	Components of breast milk	Primary/Child	Insulin resistance: - Serum insulin levels, Glucose levels, HgbA1C Obesity: - IGF

			- Body size and habitus - Body composition
Methods	Breast milk sample	Methods	Blood sample, Physical exam
Life Stage	Birth & 1-2 months	Life Stage	Birth & Years 1, 2, 5, 10, 15, 20

Important Confounders/Covariates		
Lipid profile	Increased lipid levels are associated with an increased risk of insulin resistance ¹²	
Glucokinase mutation	Glucokinase mutation is associated with increased risk of maturity onset diabetes of the young (MODY) ¹³	
Hormone levels such as cortisol, growth hormone, insulin-like growth factors	Elevated levels of these and other hormones are associated with obesity and insulin resistance in children ¹⁴	
Genetic markers for obesity	Certain genetic markers increase risk of obesity	
Parents' body mass indices	BMI and obesity are associated with certain genetic markers. ¹⁵	
Family history of diabetes and obesity	A family history of diabetes and obesity increases child's risk. 16, 15	
Lifestyle factors	Less active lifestyles would increase risk of obesity and insulin resistance. ¹⁰	
Nutrition	Poor nutritional and high caloric diet would increase risk of obesity and insulin resistance ¹⁷	
Socio-economic status and demographics	Children of lower economic status, ethnic and racial groups (particularly Native Americans, Hispanics, African Americans, and Asians) are at higher risk of obesity and insulin resistance. ¹⁶	

Population of Interest	Estimated Effect that is Detectable
All pregnant women and their offspring.	Assuming a breastfeeding prevalence of 50% (in line with estimates from NHANES III), ¹⁸ and using the power assumptions noted above, the smallest detectable relative risk for obesity would be 1.05; and for metabolic syndrome would be 1.3.

Other Design Issues		
Ethical/Burden Considerations	Blood studies, especially fasting, in younger children will require careful attention. Obtaining consent for the use of DNA may be an issue.	
Cost/Complexity of Data Collection	Addressing this hypothesis based on obesity and insulin resistance measures at later life-stages may be adversely impacted by attrition of study subjects.	

References:

¹ Parsons TJ, Power C, Logan S, Summerbell CD. Childhood predictors of adult obesity: a systematic review. Int J Obes Relat Metab Disord. 1999 Nov;23 Suppl 8:S1-107.

² Gillman MW, Rifas-Shiman SL, Camargo CA Jr, Berkey CS, Frazier AL, Rockett HR, Field AE, Colditz GA. Risk of overweight among adolescents who were breastfed as infants. JAMA. 2001 May 16;285(19):2461-7.

- ¹⁰ Sinha R, Fisch G, Teague B, Tamborlane WV, Banyas B, Allen K, Savoye M, Rieger V, Taksali S, Barbetta G, Sherwin RS, Caprio S. Prevalence of impaired glucose tolerance among children and adolescents with marked obesity. N Engl J Med. 2002 Mar 14;346(11):802-10.
- ¹¹ Colditz GA. Economic costs of obesity. American Journal of Clinical Nutrition. 55 (2): S503-S507 Suppl. S Feb 1992.
- ¹² McGarry JD: Banting Lecture 2001: Dysregulation of Fatty Acid Metabolism in the Etiology of Type 2 Diabetes. Diabetes 51:7-18, 2902
- ¹³ Kahn C.R., Vicent D., Doria A. Genetics of Non-Insulin-Dependent (Type-II) Diabetes Mellitus. Annual Review of Medicine 47: 509-531. Feb 1996.
- ¹⁴ Reinehr T., Andler W. Cortisol and Its Relation to Insulin Resistance before and after Weight Loss in Obese Children. Hormone Research 2004;62:107-112.
- ¹⁵ Maes, Hermine H.M.; Neale, Michael C.; Eaves, Lindon J. Genetic and Environmental Factors in Relative Body Weight and Human Adiposity. *Behavior Genetics*, Jul 1997; 27(4):325-51.
- ¹⁶ Fact Sheet: SEARCH for Diabetes in Youth. CDC: Department of Heath and Human Services http://www.cdc.gov/diabetes/pubs/factsheets/search.htm
- ¹⁷ Ogden CL, Flegal KM, et al. Prevalence and trends in overweight among US children and adolescents, 1999-2000. JAMA. 9 Oct 2002; 288(14): 1728-32.
- ¹⁸ Hediger ML, Overpeck MD, Kuczmarski RJ, Ruan WJ. Association between infant breastfeeding and overweight in young children. JAMA. 2001 May 16;285(19):2453-60.

³ von Kries R, Koletzko B, Sauerwald T, von Mutius E, Barnert D, Grunert V, von Voss H. Breast feeding and obesity: cross sectional study. BMJ. 1999 Jul 17;319(7203):147-50.

⁴Dietz WH. Breastfeeding may help prevent childhood overweight. JAMA. 2001 May 16;285(19):2506-7.

⁵ U.S Department of Health and Human Services, Health Resources and Services Administration, Maternal and Child Health Bureau. Child Health USA 2002. Rockville, MD, 2002.

⁶ Hedley, A.A., Ogden, C.L., Johnson, C.L., Carroll, M.D., Curtin, L.R., and Flegal, K.M. 2004. Prevalence of Overweight and Obesity Among US Children, Adolescents, and Adults, 1999-2002. JAMA 291(23): 2847-2850.

⁷ Kuczmarski RJ, Flegal KM, Campbell SM, Johnson CL. Increasing prevalence of overweight among US adults. The National Health and Nutrition Examination Surveys, 1960 to 1991. JAMA. 1994 Jul 20;272(3):205-11.

⁸ Serdula MK, Ivery D, Coates RJ, Freedman DS, Williamson DF, Byers T. Do obese children become obese adults? A review of the literature. Prev Med. 1993 Mar;22(2):167-77.

⁹ Freedman DS, Khan LK, Dietz WH, Srinivasan SR, Berenson GS. Relationship of childhood obesity to coronary heart disease risk factors in adulthood: the Bogalusa Heart Study. Pediatrics. 2001 Sep;108(3):712-8.